

DAIRYLAND BRAND IODINE TEAT DIP CONTROLLED- iodine liquid
Stearns Packaging Corporation

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.

Controlled Iodine Teat Dip

**TEAT DIPPING is an important part of a
MASTITIS CONTROL PLAN:**

1. Dip teats immediately after milking.
2. Use good milking and environmental management procedures.
3. Use properly functioning milking equipment.
4. Treat every quarter of every cow at dry off.
5. Identify and treat or manage clinical cases promptly.
6. Cull chronically infected cows.

GREAT FOR FOAMING

TEAT DIPPING IN COLD WEATHER:

When the temperature falls below 10 degrees F, or when wind chill is significant, special precautions should be taken to avoid chapped and frozen teats.

1. Teats should be dry before turning cows out into cold weather.
2. When teats are dipped after milking, allow 30 seconds contact, and blot off any excess teat dip with a single service towel.
3. Warm the teat dip to reduce drying time.
4. Provide windbreaks in outside areas for cows.
5. Monitor fresh cows with swollen udders and teats since they are more susceptible to chapped and frozen teats.

TAKE TIME

OBSERVE LABEL DIRECTIONS

DIRECTIONS FOR USE:

- DO NOT DILUTE.
- Dip teats immediately after milking as a routine part of milking procedures.
- Dip to THOROUGHLY COVER at least the lower one-third of the teat (preferably to the base of the udder), IMMEDIATELY after every milking.
- Empty and clean dip cups after each milking, or during milking, if they become contaminated.
- Do not pour used dip back into original container.
- KEEP FROM FREEZING
- Keep container closed to prevent contamination.
- Store container in an upright position.

KEEP OUT OF REACH OF CHILDREN

Warning

Causes serious eye irritation

☐Wear eye protection. Wash hands thoroughly after handling.

IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses,

if present and easy to do. Continue rinsing. If eye irritation persists: Get medical attention.

24 HOUR EMERGENCY CONTACT

1-800-255-3924

DAIRYLAND BRAND

CONTROLLED IODINE TEAT DIP

A SANITIZING TEAT DIP

An Aid in Reducing the Spread of Organisms Which May Cause Mastitis

ACTIVE INGREDIENTS:

Concentrated Iodophor, containing Nonylphenol, ethoxylated iodine complex. (provides 1/2% titratable iodine, equivalent to 5,000 ppm titratable iodine)

EMOLLIENT SYSTEM contains 5.0% glycerin.

KEEP OUT OF THE REACH OF CHILDREN

See side panel for Precautions and First-Aid Statements.

STEARNS PACKAGING CORPORATION

4200 Sycamore Ave. (53714), PO Box 3216, Madison, WI 53704

(800) 655-5008 • www.stearnspkg.com

NET CONTENTS: 1 U.S. GALLON (3.78 L)

<p>DIRECTIONS FOR USE:</p> <ul style="list-style-type: none"> DO NOT DILUTE. Dip teats immediately after milking as a routine part of milking procedures. Dip to THOROUGHLY COVER at least the lower one-third of the teat (preferably to the base of the udder), IMMEDIATELY after every milking. Empty and clean dip cups after each milking, or during milking, if they become contaminated. Do not pour used dip back into original container. KEEP FROM FREEZING. Keep container closed to prevent contamination. Store container in an upright position. <p>KEEP OUT OF REACH OF CHILDREN</p> <p>Warning Causes serious eye irritation. Wear eye protection. Wash hands thoroughly after handling. IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. If eye irritation persists, Get medical attention.</p> <p>MANTÉNGALA FUERA DEL ALCANCE DE LOS NIÑOS.</p> <p>Advertencia Provoca irritación ocular grave. Usar gafas de protección. Lávese las manos después de manipular este producto. EN CASO DE CONTACTO CON LOS OJOS: Actúe cuidadosamente con agua durante varios minutos. Quite las lentes de contacto, si las hay y resulta fácil. Siga atendiendo. Si persiste la irritación ocular. Consulte a un médico.</p>	<p>DAIRYLAND BRAND</p> <p>CONTROLLED IODINE TEAT DIP</p> <p>A SANITIZING TEAT DIP An Aid in Reducing the Spread of Organisms Which May Cause Mastitis</p> <p>ACTIVE INGREDIENTS: Concentrated iodophor, containing nonylphenol, ethoxylated iodine complex. (provides 1/2% titratable iodine, equivalent to 5,000 ppm titratable iodine) Emollient system contains 5.0% glycerin.</p> <p>KEEP OUT OF THE REACH OF CHILDREN See side panel for Precautions and First-Aid Statements.</p> <p>STEARNS PACKAGING CORPORATION 4200 Sycamore Ave. (53714), PO Box 3216, Madison, WI 53704 (800) 655-5008 • www.stearnspkg.com</p> <p>NET CONTENTS: 1 U.S. GALLON (3.78 L)</p>	<p>TEAT DIPPING IS AN IMPORTANT PART OF A MASTITIS CONTROL PLAN:</p> <ol style="list-style-type: none"> 1. Dip teats immediately after milking. 2. Use good milking and environmental management procedures. 3. Use properly functioning milking equipment. 4. Treat every quarter of every cow at dry off. 5. Identify and treat or manage clinical cases promptly. 6. Cull chronically infected cows. <p>TEAT DIPPING IN COLD WEATHER: When the temperature falls below 10°F, or when wind chill is significant, special precautions should be taken to avoid chapped and frozen teats.</p> <ol style="list-style-type: none"> 1. Teats should be dry before humping cows out into cold weather. 2. When teats are dipped after milking, allow 30 seconds contact, and blot off any excess teat dip with a single service towel. 3. Warm the teat dip to reduce drying time. 4. Provide windbreaks in outside areas for cows. 5. Monitor fresh cows with swollen udders and teats since they are more susceptible to chapped and frozen teats. <p>ST-200 LAB. 05-602015</p> <p>24 HOUR EMERGENCY CONTACT 24 horas servicio telefónico de respuesta a emergencias: 1-800-255-3924</p> <p>LOT NO:</p> <p>EXP. DATE:</p>	<p>INSTRUCCIONES POR EL USO:</p> <ul style="list-style-type: none"> NO DILUIRLO. Sumérjela los pezones inmediatamente después del ordeño como parte rutinario de procedimientos de ordeño. Sumérjala para cubrir a fondo al menos el tercio inferior de la teta (preferiblemente a la base de la ubre), inmediatamente después de cada ordeño. Vacío y limpie las tazas sumergiendo después de cada ordeño, o durante el ordeño, si continúan. No vierta los pezones utilizados volver a la recipiente de origen. NO DEJE QUE SE CONGELÉ. Mantenga el recipiente bien cerrado para evitar la contaminación. Mantenga el recipiente en una posición vertical. <p>BAÑO DE PEZONES ES UNA PARTE IMPORTANTE DE UN PLAN PARA CONTROL DE LA MASTITIS:</p> <ol style="list-style-type: none"> 1. Sumérjela los pezones inmediatamente después del ordeño, como parte rutinario de procedimientos de ordeño. 2. Utilice buenas prácticas de gestión para el establo y medio ambiente. 3. Utilice equipo de ordeño que funciona correctamente. 4. Tolle a cada cuarto de la ubre de cada vaca el dentro de fase de secado. 5. Identifique y trate o maneje casos clínicos con prontitud. 6. Entienda que las vacas que están infectadas crónicamente. <p>BAÑO DE PEZONES EN TIEMPO FRÍO: Cuando la temperatura cae por debajo de 10°F (-12°C), o cuando el factor de enfriamiento del viento es importante, tome precauciones especiales para evitar los pezones agrietados y congelados.</p> <ol style="list-style-type: none"> 1. Los pezones deben estar secos antes de soltar las vacas en el frío. 2. Cuando los pezones se sumergen después del ordeño, espere 30 segundos de contacto, y posteriormente secalos el exceso de baño de pezones con una toalla desechable. 3. Caliente el baño de pezones para reducir el tiempo de secado. 4. Proporcione protección contra el viento en las áreas fuera paja las vacas. 5. Vigile las vacas frescos con ubres hinchadas y pezones ya que son más susceptibles a los pezones agrietados y congelados.
---	---	--	---

DAIRYLAND BRAND IODINE TEAT DIP CONTROLLED

iodine liquid

Product Information			
Product Type	OTC ANIMAL DRUG	Item Code (Source)	NDC:60282-2200
Route of Administration	TOPICAL		
Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
IODINE (UNII: 9679TC07X4) (IODINE - UNII:9679TC07X4)	IODINE	5 g in 1 L	

Inactive Ingredients

Ingredient Name	Strength
GLYCERIN (UNII: PDC6A3C0OX)	50 g in 1 L

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:60282-2200-1	3.78 L in 1 BOTTLE, PLASTIC		
2	NDC:60282-2200-2	18.93 L in 1 PAIL		
3	NDC:60282-2200-3	56.78 L in 1 DRUM		
4	NDC:60282-2200-4	208.2 L in 1 DRUM		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		01/02/1982	

Labeler - Stearns Packaging Corporation (006069256)**Registrant** - Stearns Packaging Corporation (006069256)**Establishment**

Name	Address	ID/FEI	Business Operations
Stearns Packaging Corporation		006069256	manufacture, api manufacture

Revised: 3/2020

Stearns Packaging Corporation